




BPPV

Benign Paroxysmal Positional Vertigo

Information Sheet
Physiotherapy Department



WHAT IS BPPV?

Benign paroxysmal positional vertigo (BPPV) is an inner ear problem that causes short periods of vertigo when your head is moved in certain positions. It occurs most commonly when lying down, turning over in bed, and looking up. Even though you are still, you may feel like you are moving, or that the room is moving around you. You may also experience nausea, vomiting, sweating, and abnormal eye movements.

WHAT CAUSES BPPV?

BPPV occurs when small crystals of calcium carbonate, commonly referred to as rocks, shift in the inner ear and/or fall into another area within the balance canals. The crystals may shift for many reasons, such as, an inner ear infection, fever, concussion, or whiplash injury. BPPV can also occur along with other conditions of the inner ear such as Meniere's disease, migraines; or, it has been suggested that it is part of the "normal" aging process. The exact cause of this is unknown in some cases.

HOW IS IT TREATED?

Most people recover following manoeuvres which can be performed by qualified healthcare staff. This treatment aims to move the crystals back into place. Occasionally, if the problem has been there for a while we need to add in other exercises or repeat the manoeuvre to allow your system to return to normal. After the treatment is complete, you may feel nauseous, dizzy, or have more trouble balancing than before. This can last for several hours but should settle if you follow the aftercare advice.

WHAT SHOULD I DO AFTER TREATMENT?

As the Manoeuvre can make you feel off balance and dizzy. Be careful when moving initially until these symptoms settle. We advise doing this at night to lessen this effect.

Once head clears return to normal activity.

WILL IT COME BACK?

You may experience a recurrence of symptoms in the future; there is no proven medication that can prevent this condition. If you experience BPPV symptoms in the future, you should ask your GP to refer you back to the vestibular clinic. The crystals may be in a different place, so your treatment may be different than before.

The website gives instructions on how to do this yourself but if you are unsure for any reason please contact a health professional.

Remember, BPPV is treatable and the manoeuvres and exercises can greatly reduce your vertigo and other symptoms associated with BPPV.

For more information on BPPV please see our website
<https://www.nhslanarkshire.scot.nhs.uk/services/physiotherapy/vestibular-physiotherapy/>



CONFIDENTIALITY AND THE USE OF PATIENT INFORMATION

NHS Lanarkshire take care to ensure your personal information is only accessible to authorised people. Our staff have a legal and contractual duty to keep personal health information secure, and confidential. In order to find out more about current data protection legislation and how we process your information, please visit the Data Protection Notice on our website at www.nhslanarkshire.scot.nhs.uk or ask a member of staff for a copy of our Data Protection Notice.

NHS Lanarkshire - for local services and the latest health news visit www.nhslanarkshire.scot.nhs.uk
NHS Lanarkshire General
Enquiry Line: 0300 30 30 243

NHS inform - The national health information service for Scotland.
www.nhsinform.co.uk
Tel No: 0800 22 44 88

If you need this information in another language or format, please e-mail: Translation.
Services@lanarkshire.scot.nhs.uk



www.careopinion.org.uk

Pub. date:	December 2023
Review date:	December 2025
Issue No:	03
Department:	Physiotherapy
Clinical Lead:	

PIL.BPPVIS.19_33632.L
23_30938