




Guidelines for Best Practice for Clean Intermittent Catheterisation

in Children and
Young People



CONTENTS

Objectives and Rationale & Introduction	3
Introduction	4
Components of the Urinary System	5
Indications for Clean Intermittent Catheterisation	6
Impaired Bladder Functions – Causes	6
Associated Bladder Problems	7
Methods of Clean Intermittent Catheterisation	8
Catheters	8
Infection Control – General	9
Infection Control – Hand Hygiene	10
Infection Control – Hand Washing Technique Checklist	11
Your 5 Moments for Hand Hygiene	12
Procedural Guideline – General	13
Procedure – Male Urethral Catheterisation	14
Procedure – Female Urethral Catheterisation	15
Procedure – Mitrofanoff Catheterisation	16
Problem Solving – Female Urethral Catheterisation	17
Problem Solving – Urine	19
Anatomy	20
Holidays and Travelling Abroad	20
Glossary	21
References	22
Contact Information	22
Further Reading	23

OBJECTIVES

The aim of this document is to offer guidance on best practice to all carers and professionals especially those in the community who provide Clean Intermittent Catheterisation (CIC).

It provides guidance from the Best Practice Statement – Urinary Catheterisation and Catheter Care (NHS Quality Improvement Scotland, June 2004) and incorporates advice from current local practice.

RATIONALE

CIC is a normal part of daily living activity for some young people with bladder problems.

This procedure has a considerable positive impact on the young person's wellbeing and personal dignity and maintains a healthy urinary tract.

INTRODUCTION

This document has been developed using evidence based practice and references are sited at the end. This is for the use of the individual young person, parents, carers, nurses and support staff, to ensure appropriate use and teaching of Clean Intermittent Catheterisation (CIC).


Many people need the support of continence products – like catheters – to help them manage their continence and maintain a healthy bladder. CIC allows many individuals to gain control of their bladder and give them the opportunity to become self caring. CIC may be a short or long term intervention.

The procedure involves the introduction of a catheter into the bladder to remove urine, once the bladder is empty the catheter is removed. CIC is performed as often as required to prevent prolonged retention of urine and to assist continence. CIC can be performed using natural or artificially created channels. CIC has a lower infection rate than an indwelling catheter and should always be used if it is clinically appropriate and a practical option for the individual.

CIC requires to be carried out at regular intervals as recommended by health care professional. Health and education support workers may therefore be required to provide assistance or carry out this procedure while the young person is at school.

It has been demonstrated that the skills required to complete this task are well within the scope of all carers for the young person including parents, health care and education support workers working in the home or the community.

Components of the Urinary System



Indications for clean intermittent Catheterisation

(list is not inclusive)

- ❖ To ensure the bladder is emptied completely
- ❖ To irrigate the bladder
- ❖ To administer medication within the bladder.

Impaired Bladder Function – Causes

There are two types of bladder emptying problems for which CIC is indicated:

- ❖ Inability to empty bladder due to poor bladder action
- ❖ Inability to empty bladder due to obstruction

A mixture of these two can occur in the same young person. Below are some of the conditions for which CIC is recommended - this list is not inclusive.

- ❖ Congenital malformations.
- ❖ Bladder outflow obstruction – this happens when base of bladder is partially or completely blocked and reduces or prevents flow of urine into the urethra due to anatomy or stricture.

Associated Bladder Problems

- ❖ Bladder Over Activity - the bladder muscle contracts spontaneously leading to leakage of urine and in some cases backflow into the kidneys.
- ❖ Incomplete Bladder Emptying - if the bladder cannot be emptied fully, the urine that is left in the bladder is known as residual urine. Bacteria can grow quickly in this residual urine, causing a urinary tract infection.
- ❖ Urinary Tract Infections - it can be completely normal for anyone to have some bacteria in their urine. Those who perform CIC may have an even higher bacteria count in their urine. If the young person experiences symptoms such as temperature, vomiting and pain, this may be a urinary tract infection and a course of antibiotics will be prescribed. Serious recurrent and untreated urinary tract infections may cause damage to the kidneys.
- ❖ Urinary Reflux – backflow of urine from the bladder to the kidneys. Young people with reflux problems are at greater risk of urinary tract infections and kidney damage.
- ❖ Impaired Kidney Function - repeated upper urinary tract infections and prolonged urinary reflux may damage the kidneys over time. There is a risk that this may reduce the function of the kidneys long term.

Methods of Clean Intermittent Catheterisation

- ❖ Urethral - urethral catheterisation is a routine procedure performed with a sterile catheter inserted via the urethra into the bladder for the purpose of urine drainage, administering medications or irrigation of the urinary tract.
- ❖ Mitrofanoff - this is a surgical procedure to produce a stoma, an artificial channel into the bladder through the skin. A sterile catheter can be inserted through this to empty the bladder. The stoma which can look like an extra tummy button but not always. In some cases, this is combined with an operation to enlarge the young person's bladder (bladder augmentation).

Types of Catheters

(Catheters come in many types, materials and sizes)

- ❖ Flexible silicon or plastic tube.
- ❖ Lubricated with gel or require the lubrication to be activated by water.
- ❖ Normal drainage is usually achieved by using a catheter size between CH06 TO CH12.
- ❖ Correct size is the smallest possible for providing adequate drainage.
- ❖ Larger catheters can cause more leakage and damage and should only be used in special circumstances such as with an indwelling catheter insitu.

Infection Control – General


- ❖ A clean environment should be maintained at all times.
- ❖ Catheter equipment should be stored
 - in a dry, cool, dark place, out of direct sunlight.
 - unopened in manufacturer's protective boxes/wrapping to prevent damage and ensure the integrity/sterility of the product.
- ❖ Hand hygiene is appropriate for care, as hand transfer is a major factor in the current risk of infection to patients.
- ❖ Effective hand drying is carried out before catheter procedures, as friction created while drying further assists the removal of bacteria. Failure to dry hands can lead to cracked skin and breeding grounds for bacteria.
- ❖ Personal protective equipment (PPE) such as aprons and gloves should be used for all activities that have been assessed as carrying a risk of exposure to blood and body fluids/
- ❖ Single use gloves should be kept in manufacturer's box and only removed as required.
- ❖ All equipment used for these procedures should be single use wherever possible.
- ❖ All waste should be disposed of according to local policies and guidelines.

Infection Control – Hand Hygiene

- ❖ Follow principles of good hand hygiene before, during and after procedure.
- ❖ Cover all cuts and abrasions with water-proof dressings.
- ❖ Keep nails short and clean; do not wear nail varnish, artificial nails or extensions.
- ❖ Wrist-watches, stoned jewellery, bracelets or rings (other than 1 plain band) in the clinical environment. Hand jewellery will impede the performance of effective hand hygiene thus increasing the risk of infection.
- ❖ Sleeves should be worn above the elbow.

HOW TO WASH HANDS

1




Wet hands with water.

2




Apply enough soap to cover all hand surfaces.

3




Rub hands palm to palm.

4




Right palm over the back of the other hand with interlaced fingers and vice versa.

5



Palm to palm with fingers interlaced.

6




Backs of fingers to opposing palms with fingers interlocked.

7




Rotational rubbing of left thumb clasped in right palm and vice versa.

8




Rotational rubbing, backwards and forwards with clasped fingers of right hand in left palm and vice versa.

9



Rinse hands with water.

10



Dry thoroughly with towel.


11



Use elbow to turn off tap.


12

⌚ Steps 3-8 should take at least 15 seconds.



...and your hands are safe*.

Your 5 Moments for Hand Hygiene



1	BEFORE TOUCHING A PATIENT	WHEN? Clean your hands before touching a patient when approaching him/her. WHY? To protect the patient against harmful germs carried on your hands.
2	BEFORE CLEAN/ASEPTIC PROCEDURE	WHEN? Clean your hands immediately before performing a clean/aseptic procedure. WHY? To protect the patient against harmful germs, including the patient's own, from entering his/her body.
3	AFTER BODY FLUID EXPOSURE RISK	WHEN? Clean your hands immediately after an exposure risk to body fluids (and after glove removal). WHY? To protect yourself and the health-care environment from harmful patient germs.
4	AFTER TOUCHING A PATIENT	WHEN? Clean your hands after touching a patient and her/his immediate surroundings, when leaving the patient's side. WHY? To protect yourself and the health-care environment from harmful patient germs.
5	AFTER TOUCHING PATIENT SURROUNDINGS	WHEN? Clean your hands after touching any object or furniture in the patient's immediate surroundings, when leaving – even if the patient has not been touched. WHY? To protect yourself and the health-care environment from harmful patient germs.



World Health Organization

Patient Safety

A World Alliance for Safer Health Care

SAVE LIVES
Clean Your Hands

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this document. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use. WHO acknowledges the Hôpitaux Universitaires de Genève (HUG), in particular the members of the Infection Control Programme, for their active participation in developing this material.

May 2009

Procedural Guideline – General

Before, during and after procedure ensure the young person's dignity is maintained at all times, considering legal, moral, cultural and religious beliefs.

- ❖ Select dedicated location which is appropriate for performing the procedure.
- ❖ Young person has consented and is comfortable and happy for procedure to take place.
- ❖ Appropriate equipment including apron, gloves, goggles, catheter and drainage container is available.
- ❖ Ensure procedure is fully explained using appropriate language.
- ❖ Ensure the young person is in a suitable position for the procedure to be performed.
- ❖ Remove and loosen clothing where appropriate.
- ❖ Wash hands thoroughly.
- ❖ Prepare catheter, check expiry date and lubricate if necessary.
- ❖ Put on protective personal equipment eg: gloves, apron as required.
- ❖ Ensure genital area is cleaned using unfragranced soap or wipes, in downwards strokes, before insertion of catheter.

Procedure – Male Urethral Catheterisation

- ❖ Hold penis on sides, perpendicular (at 90-degree angle) to the body, straightening the urethra.
- ❖ Retract the foreskin gently where appropriate.
- ❖ Slide lubricated catheter into the opening of urethra, at tip of penis.
- ❖ Gently insert the catheter and advance it slowly towards the bladder.
- ❖ There may be resistance, do not pull catheter in and out, encourage young person to relax and breathe deeply until muscle relaxes.
- ❖ Advance catheter and massage bladder until urine starts to flow into the toilet, container or nappy.
- ❖ Advance catheter for further 1-2 cms. Hold in position until urine stops flowing.
- ❖ Remove catheter gently in small steps to ensure bladder is empty.
- ❖ If, as the catheter is being removed, urine starts to flow again, stop withdrawal until urine flow stops.
- ❖ Replace foreskin over glans of penis. Wash and dry penis, replace clothing as appropriate.
- ❖ Record time of catheterisation and amount of urine obtained if appropriate.
- ❖ Discard all waste following local policies and guidelines for clinical waste.

Procedure - Female Urethral Catheterisation

- ❖ Using fingers separate the labia and pull gently upwards to visualise the urethral meatus.
- ❖ Locate the urethra, which is below the clitoris and above the vagina.
- ❖ Slide lubricated catheter gently into the opening of urethra.
- ❖ Advance it slowly towards the bladder, guiding it upwards as if towards tummy button.
- ❖ There may be some resistance as the catheter tip reaches the bladder neck, continue to advance the catheter gently but firmly until the sphincter relaxes.
- ❖ Advance catheter until urine starts to flow into drainage container.
- ❖ Continue to advance catheter for further 1-2 cms.
- ❖ Hold in position until urine stops flowing and bladder is empty.
- ❖ If, as the catheter is being removed, urine starts to flow again, stop withdrawal until urine flow stops.
- ❖ Wash and dry genital area, replace clothing as appropriate.
- ❖ Record time of catheterisation and amount of urine obtained if appropriate.
- ❖ Discard all waste following local policies and guidelines for clinical waste.

Procedure - Mitrofanoff Catheterisation

- ❖ Remove ACE stopper if present.
- ❖ Insert the catheter gently into the mitrofanoff until you meet resistance and then give it a firm push to guide it further into the bladder.
- ❖ Allow urine to drain freely.
- ❖ When urine stops flowing, remove catheter.
- ❖ Dry stoma site if required and insert ACE Stopper (if used) and replace clothing.
- ❖ Record time of catheterisation and amount of urine obtained if appropriate.
- ❖ Discard all waste following local policies and guidelines for clinical waste.

PROBLEM SOLVING – FEMALE URETHRAL CATHETERISATION

PROBLEM	SOLUTION
GENITAL AREA RED OR INFLAMED, OR DISCHARGE PRESENT	<ul style="list-style-type: none"> ◆ Catheterise gently and with care ◆ Contact named Health Care Professional (if available) and contact parent/carer who should seek advice from GP
UNABLE TO PASS CATHETER DUE TO RESISTANCE	<ul style="list-style-type: none"> ◆ Attempt to repass new catheter ensuring that there is plenty of lubricant on catheter ◆ If still unable to catheterise contact named Health Care Professional (if available) and parent /carer
UNABLE TO OBTAIN URINE ON INSERTION OF CATHETER INTO URETHRA	<ul style="list-style-type: none"> ◆ Ensure catheter in correct position ◆ If first attempt fails, try once more ◆ If still unable to catheterise contact named Health Care Professional (if available) or parent/carer ◆ Arrange medical review
BLEEDING EVIDENT ON CATHETER INSERTION	<ul style="list-style-type: none"> ◆ A small amount of blood can be normal, check urine for urinary tract infection ◆ If amount of blood increases or continues contact named Health Care Professional (if available) and parent /carer who should seek information from GP
UNABLE TO PASS CATHETER DUE TO SPASM	<ul style="list-style-type: none"> ◆ Wait a few minutes to allow spasm to pass DO NOT FORCE CATHETER ◆ If resistance continues contact named Health Care Professional (if available) and parent/ carer

PROBLEM	SOLUTION
UNABLE TO WITHDRAW CATHETER	<ul style="list-style-type: none"> ◆ Wait a few minutes to allow spasm to pass DO NOT PULL CATHETER ◆ Wait until spasm has passed then ask child to cough and remove catheter ◆ If resistance continues contact named Health Care Professional (if available) and parent / carer
STOMA SITE RED, INFLAMED OR BLEEDING	<ul style="list-style-type: none"> ◆ Passing the catheter into the mitrofanoff can sometime cause this, but it usually stops when catheter is removed ◆ If this continues contact named Health Care Professional (if available) and contact parent /carer who should seek advice from Nurse Specialist /discharging ward at RHSC
SITE LEAKING AND SORE	<ul style="list-style-type: none"> ◆ If mucus evident, clean site and apply dry dressing to protect clothing ◆ If infection suspected contact named Health Care Professional (if available) and parent / carer to arrange for bacteriology swab to be obtained, and seek advice from GP
UNABLE TO INSERT CATHETERURINE LEAKING FROM STOMA BETWEEN CATHETERISATIONS	<ul style="list-style-type: none"> ◆ Contact named Health Care Professional (if available) and parent / carer who should seek advice from Nurse Specialist / discharging ward at RHSC
BLEEDING EVIDENT ON CATHETER INSERTION	<ul style="list-style-type: none"> ◆ A small amount of blood can be normal, check urine for urinary tract infection ◆ If amount of blood increases or continues contact named Health Care Professional (if available) and parent /carer who should seek information from GP


PROBLEM SOLVING – URINE

PROBLEM	SOLUTION
<p>PALE, STRAW COLOURED URINE. NO FOUL SMELL. NO PAIN OR DISCOMFORT.</p>	<ul style="list-style-type: none"> ◆ No action
<p>CONCENTRATED. DARK URINE, STRONG SMELLING. NO PAIN OR DISCOMFORT.</p>	<ul style="list-style-type: none"> ◆ Increase oral fluids ◆ Contact parent or carer
<p>BLOOD VISIBLE IN URINE. FOUL SMELL. PAIN AND DISCOMFORT.</p>	<ul style="list-style-type: none"> ◆ Contact named Health Care Professional (if available) or parent /carer who should seek advice from GP
<p>BLEEDING EVIDENT ON CATHETER INSERTION</p>	<ul style="list-style-type: none"> ◆ A small amount of blood can be normal, check urine for urinary tract infection ◆ f amount of blood increases or continues contact named Health Care Professional (if available) and parent /carer who should seek information from GP

ANATOMY

Male

Female



HOLIDAYS AND TRAVELLING ABROAD

Essentials for Holiday

- ◆ Order supplies for length of holiday and include extra to allow for delays.
- ◆ Arrange standby course of antibiotics from GP.
- ◆ Obtain travel letter from GP or Specialist Nurse.
- ◆ Ensure appropriate travel insurance is in place.

Essentials for Travelling Abroad

- ◆ If booking flights, ensure adequate baggage allowance for supplies.
- ◆ Consider the hygiene needs of the country you are visiting.
- ◆ Appreciate the need to carry medication and catheters in hand luggage.

GLOSSARY

ACE stopper	Small soft plug used to seal Mitrofanoff stoma
Backflow	Flow of urine in opposite direction to normal flow
Bacteria	Small organism which can cause infection
Bladder	Muscular sack where urine is stored
Bladder augmentation	Surgery to increase bladder size
Body fluids	Liquids from inside the body eg: urine, blood
Catheter	Flexible tube that can be passed into the body
Clitoris	Sexual organ situated above the opening to urethra and bladder.
Congenital	Condition present at birth
Continnence	Ability to store and release urine and faeces
Foreskin	Folds of skin that cover the tip of penis
Genital	Visible sex organs eg: penis, vagina
Glans	Sensitive area at top of penis
Hypotonic	Decreased muscle tone
Intermittent	Action which stops and starts
Irrigation	Flushing
Kidneys	The organs that remove water soluble waste products from the body by producing urine
Labia	Folds of skin around vaginal opening
Mitrofanoff	Surgical procedure to make channel through skin and into the bladder
Neurodisability	Disability affecting the nervous system
Neurogenic	A problem originating from the nervous system
Penis	Male genital organ containing the urethra
Perpendicular	At right angles
Prostate	The gland situated near the penis
Retention	Urine that is unable to be released
Retract	To draw or pull back
Sphincter	Ring like muscle
Stoma	A natural or surgically created channel
Stricture	Narrow section of urethra
Ureters	The two tubes that drain urine from kidneys to the bladder
Urethra	The tube through which urine passes out of the body
Urethral meatus	The urethral opening
Urinary tract	Produces, stores and eliminates urine
Vagina	The channel leading from genital area to womb

References

Alder Hey Children's NHS Foundation Trust. Clean Intermittent Catheterisation. 2021 European Society for Paediatric Urology Nurses. Guidelines for Intermittent Catheterisation in Children. 2016.

Great Ormond Street Hospital. Clean intermittent catheterisation. 2018.

The Royal Children's Hospital Melbourne. Teaching and supporting Clean Intermittent Catheterisation for parents and children. 2019.

Royal Manchester Children's Hospital. Clean Intermittent Catheterisation. 2021.

Seattle Children's. Clean Intermittent Catheterization Through the Urethra or a Mitrofanoff Stoma. 2019.e.

Contact Information

If you have any questions in relation to this training pack, please do not hesitate to contact the Integrated. Community Children's Nursing Team in NHS Lanarkshire.

Team Leader

INTEGRATED COMMUNITY
CHILDREN'S NURSING TEAM
NHS LANARKSHIRE
CCNTEAM@LANARKSHIRE.SCOT.
NHS.UK

PRACTICE DEVELOPMENT CENTRE
WWW.LANPDC.SCOT.NHS.UK

WISHAW GENERAL HOSPITAL
NETHERTON STREET
WISHAW, ML2 0DP

TELEPHONE: COMMUNITY
CHILDREN'S NURSE:
01698 366035.

Further Reading

1. Alencar V., Gomes C., Miranda E.P., et al. Impact of the route of clean intermittent catheterization on quality of life in children with lower urinary tract dysfunction. 2018.
2. Atchley T., Dangle P.P., Hopson B.D., et al. Age of transition to self-clean intermittent catheterization in patients with Spina Bifida. 2017
3. T. J. Atchley, P. P. Dangle, B. D. Hopson, et al. Age and factors associated with selfclean intermittent catheterization in patients with spina bifida. J PEDIATR REHABIL MED. 2018 1011 4 283-291.
4. Tariq Burki, Abdelazim Abasher, Ahlam Alshahrani, et al. Complications and patient satisfaction with urethral clean intermittent catheterization in spina bifida patients: comparing coated vs uncoated catheters. J. Pediatr. Urol. 2019 Dec15 6 646-650.
5. Chang P.Y. and S. Y. Ting. Vesicocutaneous fistula: a better way to conduct clean intermittent catheterization. 2020.
6. Dangle P.P., Atchley T., Hopson B.D., et al. Factors associated with transition from caregiver to self clean intermittent catheterization in Spina Bifida. 2017
7. Marla Andreia Garcia de Avila, Thais Rabello, Maria Paula Bortoleti de Araujo, et al. Development and Validation of an Age-Appropriate Website for Children Requiring Clean Intermittent Catheterization. Rehabil. Nurs. 2021 Mar-Apr 0146 2 65-72.
8. William DeFoor, Pramod Reddy, Melissa Reed, et al. Results of a prospective randomized control trial comparing hydrophilic to uncoated catheters in children with neurogenic bladder. J.Pediatr.Urol. 2017 Aug13 4 373.e1-373.e5.
9. Mostafa M. Elghandour, Osama El Naggat and Mohammed Abdel-Latif. Vesicocutaneous fistula: A modified technique for intermittent catheterization in children with impaired bladder function. J. Pediatr. Urol. 2021 0817 4 524.e1-524.e6.
10. Elzeneini W., Waly R., Marshall D. and A. Bailie. Early start of clean intermittent catheterization versus expectant management in children with spina bifida. J. Pediatr. Surg. 2019 February 201954 2 322-325.
11. L. Harper, O. Dunand and E. Dobremez. The fallow mitrofanoff. J. Pediatr. Urol. 2019 May15 3 261.e1-261.e4.
12. Cenita Sam, Cindrel Jagadeesan, Sudipta Sen, Pavai Arunachalam, B. Appalaraju and Praseetha Das. Urinary tract infection in pediatric patients on clean intermittent catheterization via a mitrofanoff port with reused catheters – Any association with catheter sterility? J INDIAN ASSOC PEDIATR SURG. 2020 Mar25 2 91-95.
13. Stein R., Bogaert G., Dogan H.S., et al. EAU/ESPU guidelines on the management of neurogenic bladder in children and adolescent part II operative management. Neurourol. Urodyn. 2020 01 Feb 202039 2 498-506.
14. Nordin Zakaria, Ariffin Nasir, Hans Van Rostenberghe and Mohamad Ikram Ilias. Sterile Water Versus Chlorhexidine as Cleaning Agent for Clean Intermittent Catheterization in Children: A Randomized Controlled Trial. Int. Med. J. 2020 1227 6 666-669.

CONFIDENTIALITY AND THE USE OF PATIENT INFORMATION

NHS Lanarkshire take care to ensure your personal information is only accessible to authorised people. Our staff have a legal and contractual duty to keep personal health information secure, and confidential. In order to find out more about current data protection legislation and how we process your information, please visit the Data Protection Notice on our website at www.nhslanarkshire.scot.nhs.uk or ask a member of staff for a copy of our Data Protection Notice.



NHS Lanarkshire - for local services and the latest health news visit www.nhslanarkshire.scot.nhs.uk
NHS Lanarkshire General
Enquiry Line: 0300 30 30 243

NHS inform - The national health information service for Scotland.
www.nhsinform.co.uk
Tel No: 0800 22 44 88

If you need this information in another language or format, please e-mail: Translation.
Services@lanarkshire.scot.nhs.uk



www.careopinion.org.uk

Pub. date:	December 2021
Review date:	December 2023
Issue No:	01
Department:	Paediatrics
Clinical lead:	S. Barrie

PIL.GUIDES.21_23212.W