

Irritable Hip

Information for patients



WHAT IS THE TREATMENT?

At first you should rest your child and give them regular anti-inflammatory medicines such as ibuprofen. If your child is unable to take ibuprofen because of asthma, or because they cannot tolerate it, then paracetamol should be given. We advise that children should be completely rested at home and be kept off nursery or school. They should avoid games and exercise for a period of 5 to 7 days. After 5 to 7 days your child will be reviewed in the Emergency Department clinic.

WHAT HAPPENS THEN?

If your child is better no further treatment is required. Your child should continue to avoid sports for another 1 to 2 weeks to prevent the symptoms from returning. If your child has only recently started walking they may have a slightly altered walk for a short period of time.

If your child's symptoms are persisting they may have to have blood tests and/or x-rays (if they haven't had these already). We may request an ultrasound scan of their hip.

In a very few cases children may have to be referred to see a specialist if the symptoms are failing to settle.

WHAT DO I NEED TO DO?

Give your child regular anti-inflammatories, for example ibuprofen (or paracetamol if they cannot have ibuprofen). Rest your child.

Your child needs to return to hospital if:

- ❖ They develop a high temperature.
- ❖ They develop worsening pain.

DOES THIS CONDITION COME BACK AGAIN?

Irritable hip can re-occur but this is very rare. Irritable hip is unlikely to cause long term problems.

Emergency Department

Hairmyres	01355 584716
Monklands	01236 712191
Wishaw General	01698 366630
NHS 24	08454 242424



www.careopinion.org.uk

NHS Lanarkshire - for local services and the latest health news visit www.nhslanarkshire.scot.nhs.uk

NHS Lanarkshire General Enquiry Line: 08453 130 130

If you need this information in another language or format, please e-mail Translation.Services@lanarkshire.scot.nhs.uk

Pub. date:	January 2022
Review date:	January 2024
Issue No:	04
Clinical Lead:	